

Supplementary Table. Percentage of children introduced to complementary foods before age 4 months by state among children 1–5 years – National Survey of Children's Health, United States, 2016–2018.

	Total no.	Percentage introduced early (%)*	95% CI (%)*
Alabama	432	38.7	(32.8, 44.8)
Alaska	475	28.9	(24.0, 34.4)
Arizona	422	28.8	(23.2, 35.1)
Arkansas	452	39.3	(32.6, 46.3)
California	417	23.0	(18.1, 28.8)
Colorado	486	26.6	(21.5, 32.4)
Connecticut	443	31.5	(25.8, 37.9)
Delaware	451	33.9	(28.0, 40.3)
District of Columbia	699	24.9	(20.2, 30.3)
Florida	417	35.2	(28.5, 42.5)
Georgia	398	37.9	(31.6, 44.6)
Hawaii	488	30.9	(25.9, 36.4)
Idaho	492	32.6	(27.7, 38.0)
Illinois	502	31.4	(25.7, 37.6)
Indiana	437	34.8	(29.2, 41.0)
Iowa	469	29.0	(23.7, 35.1)
Kansas	503	30.3	(24.6, 36.6)
Kentucky	459	35.8	(30.1, 42.0)
Louisiana	391	35.9	(29.9, 42.4)
Maine	448	32.2	(26.6, 38.4)
Maryland	454	35.0	(29.0, 41.4)
Massachusetts	500	33.6	(27.7, 39.9)
Michigan	486	36.3	(30.3, 42.9)
Minnesota	595	26.1	(21.0, 31.9)
Mississippi	416	49.0	(42.5, 55.5)
Missouri	481	28.5	(23.4, 34.3)
Montana	481	32.8	(27.0, 39.2)
Nebraska	550	31.2	(25.7, 37.3)
Nevada	427	35.1	(28.8, 41.9)
New Hampshire	426	30.9	(25.3, 37.1)
New Jersey	435	34.4	(28.4, 41.0)
New Mexico	382	18.0	(13.9, 23.1)
New York	446	37.3	(30.9, 44.2)
North Carolina	442	34.9	(28.4, 41.9)
North Dakota	525	28.1	(22.8, 34.2)
Ohio	474	38.4	(32.3, 44.9)
Oklahoma	458	32.6	(26.9, 38.9)
Oregon	491	22.6	(16.9, 29.5)
Pennsylvania	477	29.5	(24.1, 35.5)
Rhode Island	440	32.5	(27.1, 38.5)
South Carolina	475	29.1	(23.5, 35.3)
South Dakota	527	28.8	(23.7, 34.5)
Tennessee	444	33.3	(27.7, 39.4)

Texas	413	33.1	(27.1, 39.7)
Utah	552	36.7	(31.6, 42.2)
Vermont	478	28.0	(22.6, 34.1)
Virginia	473	32.2	(26.6, 38.3)
Washington	517	25.1	(20.1, 30.8)
West Virginia	401	36.3	(29.9, 43.2)
Wisconsin	514	27.7	(22.4, 33.7)
Wyoming	466	30.3	(25.4, 35.7)

Abbreviations: CI = confidence interval.

*Percentages are weighted to account for complex survey design.